

ADD BLUEGOLD 30-40ML/KG OF FEED TO PREVENT
LEACHING OF ALL SUPPLEMENTS IN WATER.
BETTER STABILITY AND RESILIENCE OCCURS TO PELLET.



Pack 17Lt

**THE BEST PRODUCTS FOR YOU AT FACTORY PRICE.
SOLD EXCLUSIVELY AT FACTORY DEPOT .**

FACTORY DEPOTS:

AKIVIDU - 984888789	AKIVIDU - 984888789	AKIVIDU - 984888789	AKIVIDU - 984888789	AKIVIDU - 984888789	AKIVIDU - 984888789
AKIVIDU - 984888789	AKIVIDU - 984888789	AKIVIDU - 984888789	AKIVIDU - 984888789	AKIVIDU - 984888789	AKIVIDU - 984888789


FACTORY ADDRESS:
BLUEWEIGHT BIOSCIENCES PVT LTD.
UNIT 1,13-39, BESIDES MODEL DIARY FACTORY
NIDAMANURU, VIJAYAWADA (R), INDIA AP 521104

R&D WING
BLUEWEIGHT BIOTECH LLP,
#101-45,84TH AVE, EDMONTON,
ALBERTA, CANADA. T6E 2B8

WWW.BLUEWEIGHT.COM
INFO@BLUEWEIGHT.COM
080-19504508, 8074740357



IMMUNITY PROTOCOL



**WEAK SHRIMP
DIE**



**STRONG SHRIMP
SURVIVE**



**BUILDING INNER STRENGTH AND IMMUNITY
GIVES THE HIGHEST PROBABILITY TO SUCCESS**



WEAK SHRIMP



Turns to **BRIGHT & HEALTHY SHRIMP**



DOSE
2-3/KG
OF FEED
500G PACK

- Stimulates Immune Response
- Increases resistance to Unfavourable water conditions.
- Increases Hemocyte cell count to resist disease outbreak
- Its must from the beginning to protect the value of crop
- This does benefit the shrimp in the morning, as much benefit as lemon honey water do to human.

1ST MEAL



2ND MEAL

- Builds Stamina
- Increases weight & Length of animals
- Growth Promoter
- Reduces FCR & DOC with Vitamins
- Immunostimulants and Astaxanthin



DOSE
10G/10KG
OF FEED
1KG PACK




increases Hepatopancreas Health


Removes Vibrio Pathogens inside the Hepatopancreas

Contains Allicin 12% which is working Enzyme in Garlic. In Fresh Garlic, allicin is ~0.4%, but Zyme-B contains it 12%

HEPATOPANCREAS VIBRIO LEVELS.



BEFORE



AFTER

3RD MEAL



4TH MEAL

Reduces Vibrio Pathogens in Gut. Maintains Gut pH for proper digestion.

Should be kept as last meal of the day. Act all Night.

Increases digestion of Nutrients and muscle production

Prevents the incidence of white feces if kept from beginning

DOSE
10G/10KG
OF FEED
1KG PACK



DOSE
5G/5KG
OF FEED
500G PACK